

Analytical approaches to the calculation of $f_+^{K^0\pi^-}(0)$ *

J. Portolés†

*Instituto de Física Corpuscular, IFIC,
CSIC-Universitat de València,
Apt. Correus 22085, E-46071 València, Spain*

$K_{\ell 3}$ decays constitute, at present, the golden modes to extract $|V_{us}|$ from experimental data. Its incertitude is dominated by the theoretical error in the determination of the vector form factor at zero transfer of momentum. I review the most relevant analytical approaches for the calculation of this parameter.

I. INTRODUCTION

Semileptonic kaon decays have the potential to provide information on the V_{us} CKM matrix element. In principle $K_{\ell 2}$ decays offer the most simple setting, but the fact that these are driven by the axial-vector QCD current, hence the kaon decay constant, diminish their capability: F_K is not given by symmetry and has to be input from theory, e.g. lattice. Something analogous happens in $K_{\ell 3}$ processes where another parameter guided by strong interactions, i.e. the vector form factor at zero transfer of momentum, has to be determined. However several fortunate features happen together in the later decays that convert them in an excellent system to extract information on $|V_{us}|$: i) Only the QCD vector current contributes to $K_{\ell 3}$ decays. Due to the conservation of this current in the $SU(N)_F$ limit (CVC), the normalization at zero transfer of momentum is known in this limit; ii) Corrections to the above result only start at second order in the symmetry breaking parameter, i.e. $\mathcal{O}[(m_s - m_d)^2]$ for three flavours [1].

The form factors of the vector current in $K_{\ell 3}$ decays are defined by :

$$\langle \pi^-(p_\pi) | \bar{s} \gamma_\mu u | K^0(p_K) \rangle = \quad (1)$$

$$f_+^{K^0\pi^-}(t) \left[(p_K + p_\pi)_\mu - \frac{\Delta_{K\pi}}{t} q_\mu \right] + f_0^{K^0\pi^-}(t) \frac{\Delta_{K\pi}}{t} q_\mu,$$

where $q_\mu = (p_K - p_\pi)_\mu$, $t = q^2$ and $\Delta_{K\pi} = M_K^2 - M_\pi^2$. The functions $f_+^{K^0\pi^-}(t)$ and $f_0^{K^0\pi^-}(t)$ are known as the vector and scalar form factors, respectively. The fact that $f_+^{K^0\pi^-}(0) = f_0^{K^0\pi^-}(0)$ allows us to write $f_{+,0}^{K^0\pi^-}(t) = f_+^{K^0\pi^-}(0) \left[1 + \lambda'_{+,0} \frac{t}{M_{\pi^+}^2} + \dots \right]$, and then factorize $f_+^{K^0\pi^-}(0)$ in the partial width as :

$$\Gamma(K_{\ell 3[\gamma]}) \sim |V_{us} f_+^{K^0\pi^-}(0)|^2 I^{K\ell}(\lambda'_{+,0}, \dots) \quad (2)$$

$$\times [1 + \Delta_{SU(2)} + \Delta_{EM}] ,$$

where $\Delta_{SU(2)}$ and Δ_{EM} embody isospin breaking effects of strong and electromagnetic origin, respectively [2]. As can be seen from Eq. (2), the determination of $|V_{us}|$ from $K_{\ell 3}$ decays relies essentially in our ability to ascertain an accurate value for $f_+^{K^0\pi^-}(0)$.

We have some basic model-independent knowledge on $f_+^{K^0\pi^-}(0)$. In the $SU(3)$ limit, CVC establishes that $f_+^{K^0\pi^-}(0)|_{SU(3)} = 1$. Moreover, a known sum rule [3] asserts that $f_+^{K^0\pi^-}(0) < 1$ and, consequently, flavour breaking corrections given by the Ademollo-Gatto theorem [1] should be negative. Given this setting our goal in order to provide a close determination of $f_+^{K^0\pi^-}(0)$ is to evaluate $SU(3)$ breaking corrections to $K_{\ell 3}$ decays. The appropriate framework to perform this task is Chiral Perturbation Theory (χ PT) [4, 5].

II. $f_+^{K^0\pi^-}(0)$ UP TO $\mathcal{O}(p^6)$ IN χ PT

$SU(3)$ breaking contributions span the deviation of $f_+^{K^0\pi^-}(0)$ from unity :

$$f_+^{K^0\pi^-}(0) - 1 = f_{p^4} + f_{p^6} + \dots, \quad (3)$$

where f_{p^n} is of $\mathcal{O}(p^n)$ in the chiral expansion. The $\mathcal{O}(p^4)$ correction amounts to a pure one-loop calculation (there is no local contribution from counterterms) and then it is essentially free from uncertainties, giving $f_{p^4} = -0.0227$ [6, 7]. At $\mathcal{O}(p^6)$ the situation is more complex. The first estimate was obtained by relating $f_+^{K^0\pi^-}(0)$ to the matrix element of the vector charge between a kaon and a pion in the infinite-momentum limit [6]. This procedure understates chiral log contributions and gives :

$$f_{p^6}^{LR} = -0.016(8), \quad (4)$$

that essentially coincides with the result provided by the light-front quark model [8].

Within χ PT all kind of possible contributions appear [9, 10] :

$$f_{p^6} = f_{p^6}^{2\text{-loops}}(\mu) + f_{p^6}^{L_i \times \text{loop}}(\mu) + f_{p^6}^{\text{tree}}(\mu) . \quad (5)$$

Loop terms can be evaluated with small uncertainty [10]

*Talk given at the 4th International Workshop on the CKM Unitarity Triangle, CKM2006, 12th-16th December 2006, Nagoya (Japan). Report IFIC/07-14.

†Electronic address: Jorge.Portoles@ific.uv.es

obtaining :

$$\begin{aligned} f_{p^6}^{2\text{-loops}}(M_\rho) &= 0.0113, \\ f_{p^6}^{L_i \times \text{loop}}(M_\rho) &= -0.0020(5), \end{aligned} \quad (6)$$

while the tree-level part is, a priori, unknown and can be expressed in terms of chiral low-energy couplings (LECs) of the chiral Lagrangian both at $\mathcal{O}(p^4)$ [11] and $\mathcal{O}(p^6)$ [12] :

$$f_{p^6}^{\text{tree}}(M_\rho) = 8 \frac{\Delta_{K\pi}^2}{F_\pi^2} \times \left[\frac{(L_5^r(M_\rho))^2}{F_\pi^2} - C_{12}^r(M_\rho) - C_{34}^r(M_\rho) \right], \quad (7)$$

where $F_\pi = 92.4 \text{ MeV}$ is the pion decay constant. While the L_i couplings are rather well known both from direct phenomenology [13] and on more theoretical grounds [14], our knowledge on the $\mathcal{O}(p^6)$ C_i LECs is still rather poor. As a consequence it is difficult to provide a determination for the $\mathcal{O}(p^6)$ contribution to $f_+^{K^0\pi^-}(0)$. In Ref. [10] the authors did combine their loop evaluation together with the Leutwyler & Roos (LR) result $f_{p^6}^{LR}$ in Eq. (4), assuming that the latter stands for the C_i contribution. As suggested in Ref. [15] it is more natural to presume that $f_{p^6}^{LR}$ indicates the full tree contribution $f_{p^6}^{\text{tree}}$. This would give :

$$f_{p^6}^{BT} = -0.007(9). \quad (8)$$

Several strategies to determine f_{p^6} have been developed : on one side lattice quenched and unquenched evaluations are now at hand [16], on the other there are theoretical tools that may provide a determination of the unknown LECs. The latter are usually encoded as analytical evaluations and will be considered in the following.

III. LECS FROM THE SCALAR $K\pi$ FORM FACTOR

The $\mathcal{O}(p^6)$ determination of the $K_{\ell 3}$ form factors shows that the scalar form factor in Eq. (1) depends on the same $\mathcal{O}(p^6)$ LECs than $f_{p^6}^{\text{tree}}$ [10, 17]. It can be written as :

$$\begin{aligned} f_0^{K^0\pi^-}(t) &= f_+^{K^0\pi^-}(0) + \overline{\Delta}(t) + \left(\frac{F_K}{F_\pi} - 1 \right) \frac{t}{\Delta_{K\pi}} \\ &+ \frac{8}{F_\pi^2} [\Sigma_{K\pi} (2C_{12}^r + C_{34}^r) t - C_{12}^r t^2], \end{aligned} \quad (9)$$

where $\Sigma_{K\pi} = M_K^2 + M_\pi^2$, $\overline{\Delta}(t)$ receives contributions from $\mathcal{O}(p^4)$ and $\mathcal{O}(p^6)$ and its only dependence in LECs involves L_i at $\mathcal{O}(p^6)$. Moreover $\overline{\Delta}(0) = 0$. Hence it is possible to extract the C_i LECs from data if a good knowledge on the slope and curvature of the scalar form factor is available :

$$\begin{aligned} 2C_{12}^r + C_{34}^r &\sim \frac{d}{dt} f_0^{K^0\pi^-}(t) \Big|_{t=0}, \\ C_{12}^r &\sim \frac{d^2}{dt^2} f_0^{K^0\pi^-}(t) \Big|_{t=0}. \end{aligned} \quad (10)$$

This procedure has been addressed in Ref. [17]. The $K\pi$ scalar form factor was reconstructed from a coupled-channel dispersion relation analysis from $K\pi$ scattering data [18] and then employed to obtain information on the LECs using Eq. (10) above. By adding $f_{p^6}^{\text{tree}}(M_\rho)$ to the chiral logs they obtain :

$$f_{p^6}^{\text{JOP}} = -0.003(11), \quad (11)$$

where most of the error arises from the determination of the C_i LECs. This tiny value comes from an almost complete cancellation between the loop and tree contributions.

The determination of the $K\pi$ scalar form factor from $K\pi$ scattering data shows that, in order to solve the coupled channel dispersion relations, two integration constants are required. The authors of Ref. [18] have input $f_0^{K^0\pi^-}(0)$ and $f_0^{K^0\pi^-}(\Delta_{K\pi})$. Accordingly their evaluation can be viewed more as a consistency check of the fact that $f_+^{K^0\pi^-}(0) = f_0^{K^0\pi^-}(0)$ than a pure determination of the C_i LECs.

IV. LECS FROM RESONANCE SATURATION

A systematic procedure to account for the resonance contributions to chiral LECs involves the construction of a Lagrangian theory in terms of $SU(3)$ (pseudo-Goldstone mesons) and $U(3)$ (heavier resonances) flavour multiplets as active degrees of freedom. This is called Resonance Chiral Theory ($\text{R}\chi\text{T}$). The method was applied in Ref. [14] to evaluate the resonance contributions to $\mathcal{O}(p^4)$ chiral LECs and has recently been extended [19] to look upon the $\mathcal{O}(p^6)$ LECs. The scheme relies in two key premises : i) The most general possible Lagrangian, including all terms consistent with assumed symmetry principles provides the most general possible S-matrix amplitude consistent with analyticity, perturbative unitarity, cluster decomposition and the specified principles of symmetry [4]; ii) It has been suggested that large- N_C QCD, embodied in an effective local Lagrangian for the study of meson dynamics, shows features that resemble, both qualitatively and quantitatively, the $N_C = 3$ case [20].

The procedure that has been devised amounts to construct the Lagrangian theory by imposing the chiral symmetry constraints on the pseudo-Goldstone mesons and unitary symmetry on the resonance fields. This guarantees the proper chiral limit. In addition short-distance dynamics is enforced on the couplings of the theory by demanding either the appropriate asymptotic behaviour for Green functions that are order parameters of the chiral symmetry breaking or the high-energy smoothing of form factors of QCD currents [21, 22].

It is well known that the phenomenological value of $\mathcal{O}(p^4)$ LECs is saturated by the lightest spectrum not included in χPT , i.e. resonances. This result seems to persist at one-loop in $\text{R}\chi\text{T}$ [23]. An analogous result

for the $\mathcal{O}(p^6)$ LECs has still not been reached though there is no hint that indicates it could happen otherwise : resonance saturation seems to be an underlying property stemming from non-perturbative QCD.

The analysis, along the lines outlined above, of the $\langle SPP \rangle$ Green function give us the following contributions of scalar and pseudoscalar resonances to the relevant LECs in Eq. (7) [15] :

$$\begin{aligned} L_5^{SP} &= \frac{F_\pi^2}{4M_S^2}, \\ C_{12}^{SP} &= -\frac{F_\pi^2}{8M_S^4}, \\ C_{34}^{SP} &= \frac{3F_\pi^2}{16M_S^4} + \frac{F_\pi^2}{16} \left(\frac{1}{M_S^2} - \frac{1}{M_P^2} \right)^2, \end{aligned} \quad (12)$$

where M_S (M_P) is the mass of the lightest multiplet of scalar (pseudoscalar) resonances. Substituting these values into f_p^{tree} in Eq. (7) we get :

$$f_p^{\text{tree}}(M_\rho) = -\frac{\Delta_{K\pi}^2}{2M_S^4} \left(1 - \frac{M_S^2}{M_P^2} \right)^2. \quad (13)$$

As can be seen in Figure 1 there is an almost complete cancellation between the two different counterterm contributions. This is a fortunate feature because the resulting dependence in the scalar resonance mass happens to be tiny. For $M_P = 1.3 \text{ GeV}$ and adding the chiral logs we get ¹ :

$$f_p^{\text{res}} = 0.007(12). \quad (14)$$

Notice that the sign of the central value is opposite to previous predictions above. However this is not significant because, as in f_p^{BT} or f_p^{JOP} , the result is compatible with zero.

V. COMPARISON

Lattice results, both quenched [24] or unquenched [16, 25] are available. In Table I we compare these with the results from analytical determinations reviewed above. Two trends are easily revealed : i) Lattice results, both quenched and unquenched, are in excellent agreement with the result by LR; ii) Analytical determinations driven by chiral logs and evaluations of the $\mathcal{O}(p^6)$ chiral LECs show a clear tension with the lattice ones, namely when LECs are determined by resonance saturation.

The LR prediction plays down $\mathcal{O}(p^6)$ chiral logs hence it may be an accurate prescription as long as these are not important. However the chiral determination (5) shows




FIG. 1: $f_p^{\text{tree}}(M_\rho)$ as a function of M_S for $M_P = 1.3 \text{ GeV}$ (solid line). We also plot the two components: the dashed line represents the term proportional to $L_5 \times L_5$, while the dotted line represents the term proportional to $-(C_{12} + C_{34})$.

TABLE I: Comparison of different predictions for $f_+^{K^0\pi^-}(0)$.

Reference	$f_+^{K^0\pi^-}(0)$
Leutwyler & Roos (LR) [6]	0.961(8)
Lattice (quenched) [24]	0.960(9)
Lattice (unquenched) [25]	0.968(11)
Lattice (unquenched) [16]	0.961(5)
χ -logs + LR (χ_{LR}) [10]	0.971(9)
χ -logs + f_p^{tree} [$K\pi$ scalar f.f.] [17]	0.974(11)
χ -logs + f_p^{tree} [resonance saturation] [15]	0.984(12)

that part of them are indeed large and positive, hence the addition of chiral logs and LR produces a cancellation that manifests in the χ_{LR} result of Table I. If the LR prediction stands for the local $\mathcal{O}(p^6)$ contribution it can be compared with $f_p^{\text{tree}}(M_\rho)$ showing a clear discrepancy : all LEC determinations conclude that the final result for $f_p^{\text{tree}}(M_\rho)$ is tiny (consistent with zero) while f_p^{LR} is comparatively large.

There is also a point to be understood with the lattice analyses. The latter unquenched result [16] agrees fully with the quenched determination from Ref. [24]. This could be expected if chiral logs were definitely small but Eq. (5) shows that this is not the case. The unquenched determination could miss the curvature provided by the chiral logs in the necessary extrapolation performed to reach the physical values of the masses and then its tiny error should be indeed larger. Finally there is another recourse. Lattice evaluations determine $\Delta f = f_+^{K^0\pi^-}(0) - 1 - f_{p^4}$, i.e. they cover all chiral $SU(3)$ symmetry breaking corrections except the first one. Though it would be most rare, if chiral corrections higher than $\mathcal{O}(p^6)$ were large these could bring an explanation to the discrepancy between the chiral and lattice

¹ For a detailed discussion on the error of this prediction please resort to Ref. [15].

results. However at present there is no expectation for an estimate of chiral $\mathcal{O}(p^8)$ contributions.

Determinations of $|V_{us}|$ from $K_{\ell 3}$ using the values of $f_+^{K^0\pi^-}(0)$ in Table I could also have a say to discern in the discussion just sketched. Taking into account the later published value $f_+^{K^0\pi^-}(0)|V_{us}| = 0.2164(4)$ [26], the LR value gives $|V_{us}| = 0.2252(19)$, χ_{LR} lowers it to $|V_{us}| = 0.2229(21)$ while *resonance saturation* brings it down to $|V_{us}| = 0.2199(27)$. All of them are away from the value obtained by the unitarity constraint $|V_{us}^{\text{unitarity}}| = 0.2275(12)$ [27] though, within errors, the latter is compatible with the LR evaluation.

VI. OUTLOOK

The study of $f_+^{K^0\pi^-}(0)$ is an all-important key to obtain an accurate value of $|V_{us}|$ from $K_{\ell 3}$ decays. Driven by $SU(3)$ breaking contributions its determination is, however, tampered by the non-perturbative nature of these corrections. As we have seen, and though the errors are still large, there is essentially no agreement between lattice evaluations and LR on one side, and chiral evaluations on the other. Several points need to be asserted

on the $\mathcal{O}(p^6)$ contributions : i) The size of the chiral logs has to be confirmed. If they stand as at present then unquenched lattice results should revise their extrapolation and error estimate. In addition the LR prediction should also be amended; ii) Local contributions have also to be analysed. Though all LEC determinations agree that their contribution is tiny, this conclusion is at odds with the LR evaluation.

In conclusion more work is needed in order to disentangle the $\mathcal{O}(p^6)$ contributions to $f_+^{K^0\pi^-}(0)$. A check of the size of the chiral logs would be very much welcome and, moreover, whether there is a failure of the resonance saturation hypothesis, employed in the determination of $\mathcal{O}(p^6)$ chiral LECs, or of the other models appointed to evaluate these local contributions, the situation deserves closer attention.

Acknowledgments

This work has been supported in part by the EU MRTN-CT-2006-035482 (FLAVIANet), by MEC (Spain) under grant FPA2004-00996 and by Generalitat Valenciana under grants ACOMP06/098 and GV05/015.

-
- [1] M. Ademollo and R. Gatto, Phys. Rev. Lett. **13** (1964) 264.
 - [2] V. Cirigliano, M. Knecht, H. Neufeld, H. Rupertsberger and P. Talavera, Eur. Phys. J. C **23** (2002) 121 [arXiv:hep-ph/0110153].
 - [3] V. De Alfaro, S. Fubini, G. Furlan and C. Rossetti, Nuovo Cim. A **62** (1969) 497.
 - [4] S. Weinberg, Physica A **96** (1979) 327;
 - [5] J. Gasser and H. Leutwyler, Annals Phys. **158** (1984) 142.
 - [6] H. Leutwyler and M. Roos, Z. Phys. C **25** (1984) 91.
 - [7] J. Gasser and H. Leutwyler, Nucl. Phys. B **250** (1985) 517.
 - [8] H. M. Choi and C. R. Ji, Phys. Rev. D **59** (1999) 034001 [arXiv:hep-ph/9807500].
 - [9] P. Post and K. Schilcher, Eur. Phys. J. C **25** (2002) 427 [arXiv:hep-ph/0112352].
 - [10] J. Bijnens and P. Talavera, Nucl. Phys. B **669** (2003) 341 [arXiv:hep-ph/0303103].
 - [11] J. Gasser and H. Leutwyler, Nucl. Phys. B **250** (1985) 465.
 - [12] J. Bijnens, G. Colangelo and G. Ecker, JHEP **9902** (1999) 020 [arXiv:hep-ph/9902437].
 - [13] G. Amorós, J. Bijnens and P. Talavera, Nucl. Phys. B **602** (2001) 87 [arXiv:hep-ph/0101127].
 - [14] G. Ecker, J. Gasser, A. Pich and E. de Rafael, Nucl. Phys. B **321** (1989) 311.
 - [15] V. Cirigliano, G. Ecker, M. Eidemüller, R. Kaiser, A. Pich and J. Portolés, JHEP **0504** (2005) 006 [arXiv:hep-ph/0503108].
 - [16] J. M. Zanotti, these proceedings; D. J. Antonio *et al.*, arXiv:hep-lat/0702026.
 - [17] M. Jamin, J. A. Oller and A. Pich, JHEP **0402** (2004) 047 [arXiv:hep-ph/0401080].
 - [18] M. Jamin, J. A. Oller and A. Pich, Nucl. Phys. B **622** (2002) 279 [arXiv:hep-ph/0110193].
 - [19] V. Cirigliano, G. Ecker, M. Eidemüller, R. Kaiser, A. Pich and J. Portolés, Nucl. Phys. B **753** (2006) 139 [arXiv:hep-ph/0603205].
 - [20] G. 't Hooft, Nucl. Phys. B **72** (1974) 461; E. Witten, Nucl. Phys. B **160** (1979) 57.
 - [21] S. Peris, M. Perrottet and E. de Rafael, JHEP **9805** (1998) 011 [arXiv:hep-ph/9805442]; M. Knecht, S. Peris, M. Perrottet and E. de Rafael, Phys. Rev. Lett. **83** (1999) 5230 [arXiv:hep-ph/9908283]; S. Peris, B. Phily and E. de Rafael, Phys. Rev. Lett. **86** (2001) 14 [arXiv:hep-ph/0007338].
 - [22] M. Knecht and A. Nyffeler, Eur. Phys. J. C **21** (2001) 659 [arXiv:hep-ph/0106034]; A. Pich, in Proceedings of the Phenomenology of Large N_C QCD, edited by R. Lebed (World Scientific, Singapore, 2002), p. 239, hep-ph/0205030; G. Amorós, S. Noguera and J. Portolés, Eur. Phys. J. C **27** (2003) 243 [arXiv:hep-ph/0109169]; P. D. Ruiz-Femenía, A. Pich and J. Portolés, JHEP **0307** (2003) 003 [arXiv:hep-ph/0306157]; J. Bijnens, E. Gamiz, E. Lipartia and J. Prades, JHEP **0304** (2003) 055 [arXiv:hep-ph/0304222]; J. Portolés and P. D. Ruiz-Femenía, Nucl. Phys. Proc. Suppl. **131** (2004) 170 [arXiv:hep-ph/0311251]; V. Cirigliano, G. Ecker, M. Eidemüller, A. Pich and J. Portolés, Phys. Lett. B **596** (2004) 96 [arXiv:hep-ph/0404004]. J. Portolés, Nucl. Phys. Proc. Suppl. **164** (2007) 292 [arXiv:hep-ph/0509279]; J. Portolés, arXiv:hep-ph/0702132.
 - [23] J. Portolés, I. Rosell and P. Ruiz-Femenía, arXiv:hep-ph/0611375.

- [24] D. Becirevic *et al.*, Nucl. Phys. B **705** (2005) 339 [arXiv:hep-ph/0403217].
- [25] C. Dawson, T. Izubuchi, T. Kaneko, S. Sasaki and A. Soni, Phys. Rev. D **74** (2006) 114502 [arXiv:hep-ph/0607162].
- [26] R. Versaci [By KLOE Collaboration], arXiv:hep-ex/0701008.
- [27] W. M. Yao *et al.* [Particle Data Group], J. Phys. G **33** (2006) 1.